

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,

please do not report the images to the
Image Problem Mailbox.

I Claim:

1. Mesogens having the following general formula:



3 wherein

4 X and Y independently are selected from the group consisting of terminal
5 functionalities and polymerizable groups, provided that, when X and Y both
6 are polymerizable groups, X and Y are other than bis- vinyl terminated
7 groups;

8 R^2 is a bulky organic group having a bulk greater than R^1 and R^3 whereby, when both
9 X and Y are polymerizable groups, said bulk is adapted to provide sufficient
10 steric hindrance to achieve a nematic state at room temperature while
11 suppressing crystallinity at room temperature, thereby providing effective
12 rheology and workability at room temperature; and
13 R^1 and R^3 are selected from groups less bulky than R^2 adapted to maintain said
14 nematic state.

1 2. The mesogens of claim 1 wherein X and Y independently are selected
2 from the group consisting of polymerizable groups.


3. The mesogens of claim ~~1~~ wherein R² is selected from the group consisting of alkyl groups having from about 1 to 6 carbon atoms and aryl groups.

1 4. The mesogens of claim 2 wherein R² is selected from the group
2 consisting of methyl groups, t-butyl groups, isopropyl groups, secondary butyl groups,
3 and phenyl groups.

1 5. The mesogens of claim 1 wherein R^2 is selected from the group
2 consisting of a methyl group and a t-butyl group.

1 6. The mesogens of claim 2 wherein R^2 is selected from the group
2 consisting of a methyl group and a t-butyl group.

1 7. Mesogens having the following general formula:



2 wherein

3 X and Y independently are selected from the group consisting of terminal
4 functionalities and polymerizable groups, said groups being independently
5 selected from the group consisting of acryloyloxy groups, methacryloyloxy
6 groups, hydroxyl groups, and acryloyloxy alkoxy groups, methacryloyloxy
7 alkoxy groups, alkoxy groups and alkoxoyl groups comprising alkyl groups
8 having from about 2 to about 12 carbon atoms, provided that, when X and Y
9 are both polymerizable groups, X and Y are other than unsubstituted bis- vinyl
10 terminated groups;

11 R^2 is a bulky organic group having a bulk greater than R^1 and R^3 whereby, when both
12 X and Y are polymerizable groups, said bulk is adapted to provide sufficient
13 steric hindrance to achieve a nematic state at room temperature while
14 suppressing crystallinity at room temperature, thereby providing effective
15 rheology and workability at room temperature; and

16 R^1 and R^3 are selected from groups less bulky than R^2 adapted to form said nematic
17 state.
18

1 8. The mesogens of claim 7 wherein said alkyl groups have from about 2
2 to about 9 carbon atoms.

1 9. The mesogens of claim 7 wherein said alkyl groups having from about
2 2 to about 6 carbon atoms.

1 10. The method of claim 7 wherein said polymerizable groups are selected
2 from the group consisting of cinnamoyloxy groups, acryloyloxy groups,
3 methacryloyloxy groups, and thioalkyloxy groups, acryloyloxy alkoxy groups, and
4 methacryloyloxy alkyloxy groups comprising an alkyl moiety having from about 2 to
5 about 12 carbon atoms, said alkyl moiety comprising CH₂ groups, wherein one or
6 more of said CH₂ groups independently can be substituted by oxygen, sulfur, or an
7 ester group; provided that at least 2 carbon atoms separate said oxygen or said ester
8 group.


1 11. The mesogens of claim 7 wherein X and Y independently are selected
2 from the group consisting of acryloyloxy alkyloxy groups and methacryloyloxy
3 alkyloxy groups.

1 12. The mesogens of claim 11 wherein n is from about 2 to about 9.

1 13. The mesogens of claim 11 wherein n is from about 2 to about 6.

1 14. The mesogens of claim 11 wherein n is 6.

1 15. Mesogens having the following general formula:



2
3 wherein

4 at least one of X or Y is a polymerizable group; and

the other of X or Y is independently selected from the group consisting of ester groups, organic acid groups, amine groups, hydroxyl groups, sulfhydryl groups, groups comprising a polymerizable unsaturated carbon-carbon bond, and spacer groups provided that, when X and Y are both polymerizable groups, X and Y are not bis-vinyl terminated groups;

R^2 is a bulky organic group having a bulk greater than R^1 and R^3 , whereby, when both X and Y are polymerizable groups, said bulk is adapted to provide sufficient steric hindrance to achieve a nematic state at room temperature while suppressing crystallinity at room temperature, thereby providing effective rheology and workability at room temperature; and R^1 and R^3 are selected from groups less bulky than R^2 adapted to form said nematic state.

16. The mesogens of claim 15 wherein said polymerizable groups are selected from the group comprising a polymerizable unsaturated carbon-carbon bond.

17. The mesogens of claim ~~15~~ wherein at least one of X or Y is selected from the group consisting of cinnamoyloxy groups.

18. The mesogens of claim ~~15~~ wherein one of X or Y is selected from the group consisting of acryloyloxy alkyloxy groups and methacryloyloxy alkyloxy groups.

19. A quantity of said mesogens of claim 1 wherein a proportion of a substituent selected from the group consisting of X, Y, and a combination thereof comprises a crystallization retardant, said proportion and said bulky organic group being effective to maintain said nematic state and to produce said effective rheology

5 and workability at room temperature.


1 20. The mesogens of claim 19 wherein said crystallization retardant
2 comprises at least one halogen atom.

1 21. The mesogens of claim 18 wherein said halogen atom is selected from
2 the group consisting of chlorine, bromine, and iodine.

1 22. The mesogens of claim 19 wherein said proportion is from about 3 to
2 about 50 mole%.

1 23. The mesogens of claim 19 wherein said proportion is from about 10 to
2 about 15 mole%.

1 24. Mesogens having the following general structure:



6 wherein

7 A is selected from the group consisting of alkyl groups and methyl-substituted alkyl

8 groups having from about 2 to about 12 carbon atoms; and

9 at least one of R and R⁴ is a polymerizable group, provided that, when R and R⁴ are
10 both polymerizable groups, R and R⁴ are not bis-vinyl terminated groups;


11 R² is a bulky organic group having a bulk greater than R¹ and R³, whereby, when both

12 R and R⁴ are polymerizable groups, said bulk is adapted to provide sufficient
13 steric hindrance to achieve a nematic state at room temperature while

14 suppressing crystallinity at room temperature, thereby providing effective
 15 rheology and workability at room temperature; and
 16 R^1 and R^3 are selected from groups less bulky than R^2 adapted to form said nematic
 17 state.

1 25. The mesogens of claim 24 wherein at least one of R and R^4 is selected
 2 from the group consisting of acryloxy groups and methacryloxy groups.

1 26. Mesogens having the following general formula:



2
 3 wherein


4 X and Y are selected from the group consisting of terminal functionalities and
 5 polymerizable groups, and at least one of X or Y comprises a bridging agent;

6 R^2 is a bulky organic group having a bulk greater than R^1 and R^3 whereby, when both

7 X and Y are polymerizable groups, said bulk is adapted to provide sufficient
 8 steric hindrance to achieve a nematic state at room temperature while
 9 suppressing crystallinity at room temperature, thereby providing effective
 10 rheology and workability at room temperature; and

11 R^1 and R^3 are selected from groups less bulky than R^2 which do not interfere with
 12 formation of said nematic state.


1 27. Mesogens having the following general formula:



4 X and Y are independently selected from the group consisting of acryloyloxy groups,
5 methacryloyloxy groups, hydroxyl groups, and acryloyloxy alkoxy groups,
6 methacryloyloxy alkoxy groups, alkoxy groups and alkoxoyl groups
7 comprising alkyl groups having from about 2 to about 12 carbon atoms,
8 wherein at least one of X or Y comprises a bridging agent;

下
口
口
口
口
下
口
下
口
口
口

1 28. Mesogens having the following general formula:



4 at least one of X or Y comprises a bridging agent; and

the other of X or Y is independently selected from groups comprising ester groups,
organic acid groups, amine groups, hydroxyl groups, sulfhydryl groups,
groups comprising a polymerizable unsaturated carbon-carbon bond, and
spacer groups;

9 R^2 is a bulky organic group having a bulk greater than R^1 and R^3 whereby, when both
 10 X and Y are polymerizable groups, said bulk is adapted to provide sufficient
 11 steric hindrance to achieve a nematic state at room temperature while
 12 suppressing crystallinity at room temperature, thereby providing effective
 13 rheology and workability at room temperature; and
 14 R^1 and R^3 are selected from groups less bulky than R^2 adapted to form said nematic
 15 state.

1 29. The mesogens of claim 26 wherein said bridging agent comprises a
 2 dicarboxoyl group comprising from about 4 to about 12 carbon atoms.

1 30. The mesogens of claim 27 wherein said bridging agent comprises a
 2 dicarboxoyl group comprising from about 4 to about 12 carbon atoms.


1 31. The mesogens of claim 28 wherein said bridging agent comprises a
 2 dicarboxoyl group comprising from about 4 to about 12 carbon atoms.

1 32. The mesogens of claim 26 wherein said bridging agent comprises an
 2 oligodialkylsiloxane comprising alkyl groups comprising from about 1 to about 3
 3 carbon atoms.

1 33. The mesogens of claim 27 wherein said bridging agent comprises an
 2 oligodialkylsiloxane comprising alkyl groups comprising from about 1 to about 3
 3 carbon atoms.


1 34. The mesogens of claim 28 wherein said bridging agent comprises an
 2 oligodialkylsiloxane comprising alkyl groups comprising from about 1 to about 3
 3 carbon atoms.

1 35. The mesogens of claim 1 wherein at least one of X or Y has the
 2 following general structure:




wherein Z is selected from the group consisting of a terminal functionality and a polymerizable group.

36. The mesogens of claim 7 wherein at least one of X or Y has the following general structure:




wherein Z is selected from the group consisting of a terminal functionality and a polymerizable group.

37. The mesogens of claim 15 wherein at least one of X or Y has the following general structure:



wherein Z is selected from the group consisting of a terminal functionality and a polymerizable group.

38. The mesogens of claim 24 wherein at least one of X or Y has the following general structure:



4

5 wherein Z is selected from the group consisting of a terminal functionality and a
6 polymerizable group.

1 39. The mesogens of claim 1 wherein R and R³ are selected from the group
2 consisting of hydrogen and a methyl group.

1 40. The mesogens of claim 7 wherein R and R³ are selected from the group
2 consisting of hydrogen and a methyl group.

1 41. The mesogens of claim 15 wherein R and R³ are selected from the
2 group consisting of hydrogen and a methyl group.

1 42. The mesogens of claim 24 wherein R and R³ are selected from the
2 group consisting of hydrogen and a methyl group.

1 43. The mesogens of claim 24 wherein said alkyl groups have from about
2 2 to about 9 carbon atoms

1 44. The mesogens of claim 24 wherein said alkyl groups have from about
2 2 to about 6 carbon atoms.

1 45. The mesogens of claim 24 wherein said alkyl groups have 6 carbon
2 atoms.


1 46. The mesogens of claim 24 wherein A is selected from the group
2 consisting of alkyl groups and methyl-substituted alkyl groups having from about 2 to
3 about 9 carbon atoms.

10056121-012302

47. The mesogens of claim 24 wherein A is selected from the group consisting of alkyl groups and methyl-substituted alkyl groups having from about 2 to about 6 carbon atoms.

48. The mesogens of claim 24 wherein A has 6 carbon atoms.

49. Mesogens having the following general formula:



R^2 is a bulky organic group having a bulk greater than R^1 and R^3 whereby, when both terminal OH groups are reacted with a polymerizable group, said bulk is adapted to provide sufficient steric hindrance to achieve a nematic state at room temperature while suppressing crystallinity at room temperature, thereby providing effective rheology and workability at room temperature; and

R^1 and R^3 are selected from groups less bulky than R^2 adapted to form said nematic state.

50. The mesogens of claim 49 wherein R^2 is selected from the group consisting of alkyl groups having from about 1 to 6 carbon atoms and aryl groups.

51. The mesogens of claim 49 wherein R^2 is selected from the group consisting of alkyl groups having from about 1 to about 4 carbon atoms and phenyl groups.

52. The mesogens of claim 49 wherein R^2 is selected from the group consisting of methyl groups, t-butyl groups, isopropyl groups, secondary butyl groups, and phenyl groups.

53. The mesogens of claim 49 wherein R^1 and R^3 are selected from the

10056121-012302

2 group consisting of hydrogen and a methyl group.

1 54. The mesogens of claim 50 wherein R and R³ are selected from the
2 group consisting of hydrogen and a methyl group.

1 55. The mesogens of claim 51 wherein R and R³ are selected from the
2 group consisting of hydrogen and a methyl group.

1 56. The mesogens of claim 52 wherein R and R³ are selected from the
2 group consisting of hydrogen and a methyl group.


1 57. The mesogens of claim 49 wherein R and R³ are hydrogen.

1 58. The mesogens of claim 50 wherein R and R³ are hydrogen.

1 59. The mesogens of claim 51 wherein R and R³ are hydrogen.

1 60. The mesogens of claim 52 wherein R and R³ are hydrogen.

1 61. Mesogens having the following general structure:



3

4 wherein

5 R^4 is an alkylene group having from about 2 to about 20 carbon atoms;

6 R^5 and R^6 are selected from the group consisting of hydrogen, halogen, and
7 bulky organic groups; and,
8 V and W independently are selected from the group consisting of terminal
9 functionalities and polymerizable groups.

1 62. The mesogens of claim 61 wherein at least one of R^5 and R^6 is a bulky
2 organic group selected from the group consisting of alkyl groups having from about 1
3 to 6 carbon atoms and aryl groups.

1 63. The mesogens of claim 61 wherein at least one of R^5 and R^6 is a bulky
2 organic group is selected from the group consisting of alkyl groups having from about
3 1 to about 4 carbon atoms and phenyl groups.

1 64. The mesogens of claim 61 wherein at least one of R^5 and R^6 is a bulky
2 organic group is selected from the group consisting of methyl groups, t-butyl groups,
3 isopropyl groups, secondary butyl groups, and phenyl groups.

1 65. The mesogens of claim 61 wherein at least one of R^5 and R^6 is selected
2 from the group consisting of methyl groups and t-butyl groups.

1 66. The mesogens of claim 61 wherein R^4 has from about 2 to about 12
2 carbon atoms.

1 67. The mesogens of claim 62 wherein R^4 has from about 2 to about 12
2 carbon atoms.

1 68. The mesogens of claim 63 wherein R^4 has from about 2 to about 12
2 carbon atoms.

1 69. The mesogens of claim 64 wherein R^4 has from about 2 to about 12
2 carbon atoms.

1 70. The mesogens of claim 61 wherein R^4 has from about 6 to about 12

10056121.012302

2 carbon atoms.

1 71. The mesogens of claim 62 wherein R^4 has from about 6 to about 12
2 carbon atoms.

1 72. The mesogens of claim 63 wherein R^4 has from about 6 to about 12
2 carbon atoms.

1 73. The mesogens of claim 64 wherein R^4 has from about 6 to about 12
2 carbon atoms.

1 74. The mesogens of claim 61 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, sulfhydryl groups, and spacer groups.

1 75. The mesogens of claim 62 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, sulfhydryl groups, and spacer groups.

1 76. The mesogens of claim 63 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, sulfhydryl groups, and spacer groups.

1 77. The mesogens of claim 64 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, sulfhydryl groups, and spacer groups.

1 78. The mesogens of claim 61 wherein said terminal functionalities are
2 hydroxyl groups.

1 79. The mesogens of claim 62 wherein said terminal functionalities are
2 hydroxyl groups.

10056121-012302

1 80. The mesogens of claim 63 wherein said terminal functionalities are
2 hydroxyl groups.

1 81. The mesogens of claim 64 wherein said terminal functionalities are
2 hydroxyl groups.

1 82. The mesogens of claim 61 wherein said polymerizable groups are
2 selected from the group consisting of alkenyl ester groups comprising a polymerizable
3 unsaturated carbon-carbon bond wherein said alkenyl group has from about 2 to about
4 12 carbon atoms.

1 83. The mesogens of claim 82 wherein said alkenyl group has from about
2 2 to about 9 carbon atoms.

1 84. The mesogens of claim 82 wherein said alkenyl group has from about
2 2 to about 6 carbon atoms.

3 85. The mesogens of claim 61 wherein V and W independently are
4 selected from the group consisting of acryloyloxy alkoxy groups and methacryloyloxy
5 alkoxy groups.

1 86. The mesogens of claim 62 wherein V and W independently are
2 selected from the group consisting of acryloyloxy alkoxy groups and methacryloyloxy
3 alkoxy groups.

1 87. The mesogens of claim 63 wherein V and W independently are
2 selected from the group consisting of acryloyloxy alkoxy groups and methacryloyloxy
3 alkoxy groups.

1 88. The mesogens of claim 64 wherein V and W independently are
2 selected from the group consisting of acryloyloxy alkoxy groups and methacryloyloxy
3 alkoxy groups.

1 89. The mesogens of claim 69 wherein V and W independently are
2 selected from the group consisting of acryloyl groups and methacryloyl groups.

1 90. The mesogens of claim 73 wherein V and W independently are
2 selected from the group consisting of acryloyloxy alkoxy groups and methacryloyloxy
3 alkoxy groups.

1 91. A composition comprising alkylenedioic bis-(4-{2-R²-4-[4-(hydroxy)-
2 benzoyloxy]-phenoxy-carbonyl}-phenyl) esters wherein R² is a selected from the
3 group consisting of alkyl groups having from about 1 to 6 carbon atoms and aryl
4 groups.

1 92. The composition of claim 91 wherein R² is selected from the group
2 consisting of alkyl groups having from about 1 to about 4 carbon atoms and phenyl
3 groups.


1 93. The composition of claim 92 wherein R² is selected from the group
2 consisting of methyl groups, t-butyl groups, isopropyl groups, secondary butyl groups,
3 and phenyl groups.

1 94. The composition of claim 91 wherein R² and R² are selected from the
2 group consisting of methyl groups and t-butyl groups.

1 95. A composition comprising decanedioic acid bis-(4-{2-tert-butyl-4-[4-
2 (2-methyl-acryloyloxy)-benzoyloxy]-phenoxy-carbonyl}-phenyl) ester.

1 96. A composition comprising a mesogen having the following general
2 structure:

3



wherein

R^5 and R^6 are selected from the group consisting of hydrogen, halogen, alkyl groups having from about 1 to 6 carbon atoms, and aryl groups; and,

V and W independently are selected from the groups comprising polymerizable groups and terminal functionalities.

97. The composition of claim 96 wherein V and W independently are selected from the group consisting of acryloyloxy groups, methacryloyloxy groups, acryloyloxy alkoxy groups and methacryloyloxy alkoxy groups.

98. The composition of claim 97 wherein R^5 and R^6 are selected from the group consisting of alkyl groups having from about 1 to about 4 carbon atoms and phenyl groups.

99. The composition of claim 97 wherein R^5 and R^6 are selected from the group consisting of methyl groups, t-butyl groups, isopropyl groups, secondary butyl groups, and phenyl groups.

100. The composition of claim 97 wherein R^5 and R^6 are selected from the

2 group consisting of methyl groups and t-butyl groups.

1 101. The composition of claim 98 wherein R^5 and R^6 are selected from the
2 group consisting of alkyl groups having from about 1 to about 4 carbon atoms and
3 phenyl groups.

1 102. The composition of claim 98 wherein R^5 and R^6 are selected from the
2 group consisting of methyl groups, t-butyl groups, isopropyl groups, secondary butyl
3 groups, and phenyl groups.

1 103. The composition of claim 98 wherein R^5 and R^6 are selected from the
2 group consisting of methyl groups and t-butyl groups.

1 104. The composition of claim 97 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, and sulfhydryl groups.

1 105. The composition of claim 97 wherein said terminal functionalities are
2 hydroxyl groups.


1 106. The composition of claim 102 wherein said terminal functionalities
2 independently are selected from the group consisting of hydroxyl groups, amino
3 groups, and sulfhydryl groups.

1 107. The composition of claim 102 wherein said terminal functionalities are
2 hydroxyl groups.

1 108. A composition comprising a mesogen having the following general
2 structure:

3

10056121-012302



$C_{66}H_{66}O_{16}$
 Exact Mass: 1114.44
 Mol. Wt.: 1115.22
 C, 71.08; H, 5.97; O, 22.95

4
5

1 109. The mesogens of claim 1 wherein said polymerizable groups are
 2 groups adapted to be polymerized by either free radical polymerization or by Michael
 3 addition.

1 110. The mesogens of claim 7 wherein said polymerizable groups are
 2 groups adapted to be polymerized by either free radical polymerization or by Michael
 3 addition.

1 111. The mesogens of claim 15 wherein said polymerizable groups are
 2 groups adapted to be polymerized by either free radical polymerization or by Michael
 3 addition.

5

10056121-012302